

Exact Derivation of Luttinger Liquid Relation in a One-Dimensional Two-Component Quantum System with Hyperbolic Interactions

Rudolf A. Römer and Bill Sutherland

Physics Department, University of Utah, Salt Lake City, Utah 84112

(February 15, 1994)

Abstract

We present an exact calculation of the Luttinger liquid relation for the one-dimensional, two-component SC model in the interaction strength range $-1 < s < 0$ by appropriately varying the limits of the integral Bethe Ansatz equations. The result is confirmed by numerical and conformal methods. By a related study of the transport properties of the SC model, we can give an exact formula for the susceptibility. Our method is applicable to a wide range of models such as, e.g., the Heisenberg-Ising chain.

72.15Nj,05.30.-d,75.30Ds

Typeset using REVTeX

The Luttinger liquid approach [1] to one-dimensional quantum models in conjunction with methods of conformal field theory [2] is extremely useful for understanding the low-energy physics of these models. The essence of the method is the realization that close to the fermi level, all excitations are essentially hydrodynamic density fluctuation plus finite-size terms conveniently described by conformal finite size formulas [3]. Furthermore, the finite-size spectrum may be given in terms of a single parameter due to the presence of the so-called Luttinger liquid relation. In most models, this parameter is the renormalized interaction strength.

Although the set of models that fall into the Luttinger liquid universality class (the $c = 1$ universality class of the Gaussian model of conformal field theory) incorporates all “classic” short-ranged 1D models such as Heisenberg-Ising and Hubbard model [4], there is no rigorous proof that *all* possible 1D models should, too. Indeed, the calculation of the central charge c in long-ranged models of the $1/r^2$ type yields contradictory results [5].

In this Letter, we will give an exact derivation of a Luttinger liquid type relation in a two-component quantum system with hyperbolic interaction defined by the Hamiltonian

$$H = -\frac{1}{2} \sum_{1 \leq j \leq N} \frac{\partial^2}{\partial x_j^2} + \sum_{1 \leq j < k \leq N} v_{jk}(x_j - x_k), \quad (1)$$

with pair potential given as

$$v_{jk}(x) = s(s+1) \left[\frac{1 + \sigma_j \sigma_k}{2 \sinh^2(x)} - \frac{1 - \sigma_j \sigma_k}{2 \cosh^2(x)} \right], \quad s > -1. \quad (2)$$

The quantum number $\sigma = \pm 1$ distinguishes the two kinds of particles. We may usefully think of it as either representing charge or spin. The model has been solved in Ref. [6] by an asymptotic Bethe Ansatz calculation and for large distances, its correlation functions have been calculated [7]. We call it the SC model for the sinh-cosh interaction. The SC model is closely related to the Hubbard and the Heisenberg-Ising (H-I) model [8] and may be viewed as a H-I fluid, with mobile spins [9].

We restrict our investigation to $-1 < s < 0$, where the system has two gapless excitations corresponding to a particle-hole and a two spin-wave continuum with Fermi velocities v_c and

v_s , respectively. Let M be the number of spin-down particles ($\sigma = -1$) and $N - M$ the number of spin-up particles for a total of N on a ring of length L . We then twist [10] the Bethe-Ansatz equations obtained in Ref. [7] for N pseudo-momenta $\mathbf{k} = (k_1, \dots, k_N)$ and M rapidities $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_M)$ with a flux Φ such that the total momentum $P(\mathbf{k}) = \sum k = 0$, i.e.,

$$Lk = 2\pi I(k) - \frac{M}{N}\Phi + \sum_{\lambda}^M \theta_1(k - \lambda) + \sum_{k'}^N \theta_0(k - k'), \quad (3a)$$

$$0 = 2\pi J(\lambda) + \Phi + \sum_{\lambda'}^M \theta_2(\lambda - \lambda') + \sum_k^N \theta_1(\lambda - k). \quad (3b)$$

The two-body phase shifts for particle-particle, particle-spin wave and spin wave-spin wave scattering, i.e., $\theta_0(k)$, $\theta_1(k)$ and $\theta_2(k)$ respectively, have been given previously [6]. The particle quantum numbers I_j and the spin-wave quantum numbers J_a are integers or half-odd integers depending on the parities of N , M as well as on the particle statistics [7].

In the thermodynamic limit, i.e., $L \rightarrow \infty$ at fixed particle and spin wave densities, $d = N/L$ and $m = M/N$, respectively, $\mathbf{k}$ and $\boldsymbol{\lambda}$ will be distributed densely around the origin. The ground state is a filled Fermi sea characterized by the distribution function $\rho(k)$ of particles and $\sigma(\lambda)$ of down-spins. so that we can replace the sums in Eq.(3) with integrals.

This yields

$$k = \frac{2\pi I(k)}{L} - \frac{M}{NL}\Phi + \int_{-C}^C \theta_1(k - \mu)\sigma(\mu)d\mu + \int_{-B}^B \theta_0(k - h)\rho(h)dh, \quad (4a)$$

$$0 = \frac{2\pi J(\lambda)}{L} + \frac{\Phi}{L} + \int_{-C}^C \theta_2(\lambda - \mu)\sigma(\mu)d\mu + \int_{-B}^B \theta_1(\lambda - h)\rho(h)dh, \quad (4b)$$

and the values of B and C are fixed by the following equations:

$$\int_{-B}^B \rho(k)dk = d, \quad (5)$$

$$\int_{-C}^C \sigma(\lambda)d\lambda = md \equiv (d - y)/2. \quad (6)$$

We rewrite these equations using the operator notation of [8]: $\langle k|\Theta_i|h\rangle \equiv \theta_i(k - h)$, $\langle k|\rho\rangle \equiv \rho(k)$, etc.. B and C are defined as projection operators to yield the finite limits of the integration. In this notation, Eq.(4) is

$$k = \frac{2\pi I(k)}{L} - \frac{M}{NL}\Phi + \Theta_1 C\sigma + \Theta_0 B\rho, \quad (7a)$$

$$0 = \frac{2\pi J(\lambda)}{L} + \frac{\Phi}{L} + \Theta_2 C\sigma + \Theta_1 B\rho. \quad (7b)$$

Let us take the derivative of Eq.(4) w.r.t. k . This then yields a set of integral equations for $\rho(k)$ and $\sigma(\lambda)$ (Ref. [7], Eq.(10)). In terms of the operator notation we have

$$1/2\pi = \rho + K_1 C\sigma + K_0 B\rho, \quad (8a)$$

$$0 = \sigma + K_2 C\sigma + K_1 B\rho, \quad (8b)$$

where the new kernels are defined as $\langle k|K_i|h\rangle \equiv \frac{1}{2\pi}\theta'_i(k-h)$.

The neutral sector of the SC model is defined as $M = N/2$, i.e. $C = \mathbb{1}$ and $y = 0$. We now wish to calculate the effect of a small disturbance $y \simeq 0$ ($C \lesssim \mathbb{1}$) and $\Phi \simeq 0$ on the system. Eq.(8b) may be written as

$$\sigma + K_2 C\sigma = -K_1 B\rho, \quad (9)$$

$$\sigma = -(1 + J_2)K_1 B\rho + (1 + J_2)K_2(1 - C)\sigma, \quad (10)$$

with J_i the resolvent of K_i . Note that as pointed out in Ref. [9], the phase-shifts θ_1 and θ_2 are essentially the phase shift of the H-I model after the identification of $|s| = \mu/\pi$. Therefore the existence of the resolvents J_1 and J_2 in the regime $-1 < s < 0$ has been proved in Ref. [8].

Let σ_0 be a solution of Eq.(8b) with $C = \mathbb{1}$, i.e., $\sigma_0 = -(1 + J_2)K_1 B\rho$. Then we may also write $\sigma + J_2(1 - C)\sigma = \sigma_0$. We substitute this back into Eq.(4a) and have

$$k = \frac{2\pi I(k)}{L} + [\Theta_0 - \Theta_1(1 + J_2)K_1] B\rho - \frac{M}{NL}\Phi - \Theta_1(1 + J_2)(1 - C)\sigma. \quad (11)$$

Defining $\Theta \equiv [\Theta_0 - \Theta_1(1 + J_2)K_1]$ and $g \equiv -\frac{M}{NL}\Phi - \Theta_1(1 + J_2)(1 - C)\sigma$ this yields

$$k = \frac{2\pi I(k)}{L} + \Theta B\rho + g. \quad (12)$$

We see that we have succeed in rewriting Eq.(3a) as the equation at $y = 0$, $\Phi = 0$ plus a small perturbation term g .

Let us briefly discuss how to handle the effect of this disturbance for additive quantities such as the energy. The net result of the presence of g will be a shift in the pseudo-momenta $k = k_0 + \delta k$, with δk of order $1/L$. Then, expanding Eq.(12), eliminating terms of order 1 with the help of the unperturbed equation and defining $\gamma = \delta k \rho$, we have $\gamma + KB\gamma = g/2\pi$. Introducing the resolvent $(1 + J)B(1 + K) = (1 + K)B(1 + J) = \mathbb{1}$, we can solve for γ , i.e., $\gamma = (1 + J)Bg/2\pi$. Let $E = \frac{1}{2} \sum^N k^2$ be the energy. Then the perturbed energy to first order will be $E = \sum E(k_0 + \delta k) \simeq \sum E(k_0) + \sum E'(k_0)\delta k$ and we calculate the difference due to the presence of the perturbation g as

$$\begin{aligned} \Delta E/L &\equiv (E(k) - E(k_0))/L \simeq BE'\delta k\rho \\ &= \int_{-B}^B E'(k)\gamma(k)dk = \gamma^+ BE' \\ &= g^+ B(1 + J)BE'/2\pi = g^+ B(1 + J)Bk/2\pi. \end{aligned} \quad (13)$$

The excitation spectrum of Bethe Ansatz models has been expressed by Yang and Yang [11] in terms of the solution to an integral equation. For the SC model the corresponding integral equation is

$$\epsilon + KB\epsilon = k^2/2 - (\text{chemical potential}). \quad (14)$$

The chemical potential is chosen such that $\epsilon(\pm B) = 0$ and thus we may take the derivative of the above equation, perform a partial integration and have $\epsilon' + KB\epsilon' = k$ or $\epsilon' = (1 + J)Bk$. Inserting this into Eq.(13) and performing another partial integration, we have the final result

$$\Delta E/L = -\epsilon^+ Bg'/2\pi = \epsilon^+ BK_1(1 + J_2)(1 - C)\sigma. \quad (15)$$

In Eq.(3b), $J(\lambda)$ was defined for $\lambda \in [-C, C]$ only. However, the equations also work outside, i.e., $0 = 2\pi J(\infty) + \Phi + (M - 1)\theta_2(\infty) + N\theta_1(\infty) = 2\pi J(\infty) + \Phi + (M - 1)\pi(1 + 2s) - N\pi(1 + s) \simeq 2\pi J(\infty) + \Phi + L[md\pi(1 + 2s) - d\pi(1 + s)]$. Thus we have

$$\begin{aligned}\int_{C_1}^{\infty} \sigma(\lambda) d\lambda &= [J(\infty) - J(C_1)]/L \\ &\simeq [\pi y(1+s) - (\Phi/L)]/2\pi,\end{aligned}\tag{16}$$

since $J(C_1) = (M-1)/2 \simeq Lmd$. The same idea works for the lower limit.

Let us now return to Eq.(9). We make the identifications $\alpha = \pi\lambda$ and $\mu = \pi|s|$. Then the equation reads as

$$\sigma(\alpha) + \frac{1}{2\pi} \int_{-\pi C_2}^{\pi C_1} \frac{\sin(2\mu)}{\cosh(\alpha - \beta) - \cos(2\mu)} \sigma(\beta) d\beta = \frac{1}{2} \int_{-B_2}^{B_1} \frac{\sin(\mu)}{\cosh(\alpha - \pi k) - \cos(\mu)} \rho(k) dk.\tag{17}$$

If we compare this with Yang and Yang's [8] integral equation $R + \mathbf{K}R = \xi$, we see that we can identify $\xi = -K_1(\alpha)/2$, $\mathbf{K} = K_2(\alpha)$ and write

$$\sigma(\lambda) = \frac{1}{2} \int_{-B_2}^{B_1} R(\pi\lambda - \pi k) \rho(k) dk.\tag{18}$$

Furthermore, we have $\sigma_0 = R_0 B \rho / 2 \simeq R_0 B \rho_0 / 2$. Again from Yang and Yang, we can then compute the asymptotic behavior of σ_0 , i.e.,

$$\begin{aligned}\sigma_0 &\xrightarrow{\alpha \rightarrow \infty} \frac{\pi}{\mu} e^{-\frac{\pi\alpha}{2\mu}} \frac{1}{2} \int_{-B_2}^{B_1} e^{\frac{\pi^2 k}{2\mu}} \rho_0(k) dk \\ &\simeq R_0(\alpha) a / 2,\end{aligned}\tag{19}$$

where a is defined as the integral. Thus we write $\sigma(\lambda) \simeq a R(\pi\lambda)/2$. Knowing the asymptotics of $R(\alpha)$ from Yang and Yang [8] and using their notation, we can compute

$$\begin{aligned}\int_{C_1}^{\infty} \sigma(\lambda) d\lambda &= \frac{a}{2\pi} \int_{\pi C_1}^{\infty} R(\alpha) d\alpha \\ &= \frac{a\zeta_1}{2\mu} \tilde{T}(0),\end{aligned}\tag{20}$$

with $\zeta_1 \equiv e^{-\pi^2 C_1 / 2\mu} \ll 1$. The same calculation may be done for the lower limit and comparison with Eq.(16) yields

$$\zeta_{1,2} = \frac{\mu}{\pi a \tilde{T}(0)} [y(\pi - \mu) \mp (\Phi/L)].\tag{21}$$

Using Eq.(15), we may then express the corrections to the energy as

$$\begin{aligned}
\Delta E/L &= -\frac{1}{2}\epsilon^+ B R_0(1-C)\sigma \\
&= -\frac{1}{2} \int_B \epsilon(k) \left[\int_{1-C} R_0(\pi\lambda - \pi k) \sigma(\lambda) d\lambda \right] dk \\
&= -\frac{\pi a}{4\mu} \int_B \epsilon(k) \left[\int_{1-C} e^{-(\pi^2\lambda - \pi^2 k)/2\mu} R(\pi\lambda) d\lambda \right] dk.
\end{aligned} \tag{22}$$

We now again use the identification $\alpha = \pi\lambda$ and then shift the integration boundaries from $\int_{C_1}^\infty$ ($\int_{-\infty}^{C_2}$) to $\int_0^\infty$ ($\int_{-\infty}^0$). Furthermore, we treat the contributions from the upper and lower limits as independent. Then we are left with the Wiener-Hopf type problem of Yang and Yang [8] and we may conveniently use their results, i.e.,

$$\Delta E/L = -\frac{a}{4} \int_B \epsilon(k) e^{\pi^2 k/2\mu} dk \frac{1}{\pi} \left[(\pi\zeta_1/\mu)^2 + (\pi\zeta_2/\mu)^2 \right] \tilde{T}(i\pi/2\mu). \tag{23}$$

The k integration may be rewritten by a partial integration

$$-\int_B \epsilon(k) e^{\pi^2 k/2\mu} = \frac{2\mu}{\pi^2} \int_B \epsilon'(k) e^{\pi^2 k/2\mu} \equiv \frac{2\mu}{\pi^2} b, \tag{24}$$

such that $\Delta E/L = \frac{ab}{2\pi\mu} \tilde{T}(i\pi/2\mu) [\zeta_1^2 + \zeta_2^2]$. Using Eq.(21), we then have

$$\Delta E/L = \frac{\tilde{T}(i\pi/2\mu)}{\tilde{T}^2(0)} \frac{b}{a} \frac{\mu}{\pi^3} \left[(\Phi/L)^2 + (\pi - \mu)^2 y^2 \right]. \tag{25}$$

Again from Yang and Yang, we know $\frac{\tilde{T}(i\pi/2\mu)}{\tilde{T}^2(0)} = \frac{\pi^2}{8\mu(\pi-\mu)}$ and furthermore $b/a = 2\pi v_s$. This gives the final result

$$\Delta E/L = \frac{v_s(\pi - \mu)}{4} \left[y^2 + \frac{(\Phi/L)^2}{(\pi - \mu)^2} \right]. \tag{26}$$

Defining the stiffness D and the susceptibility χ as $\Delta E/L = \frac{1}{2} [\chi^{-1}(y/2)^2 + D(\Phi/L)^2]$, we therefore have the desired relation

$$D\chi^{-1} = v_s^2. \tag{27}$$

As we have shown in Ref. [9], this relation may be also derived from conformal finite size formulas using the dressed charge matrix as calculated by Wiener-Hopf techniques [7,8]. Furthermore, an analogous calculation yields the Luttinger relation for the H-I model and the method may be easily extended to other models. However, we made good use of the seminal

papers of Yang and Yang [8,11] in the present exposition and correspondingly useful results do not exist for most other models. We remark that the essential idea for our calculation is already contained in these papers, and also in Haldane's "Luttinger" papers [1].

In Ref. [9], we have also shown how to relate stiffness and susceptibility by threading the system with a flux of $\Phi = 2\pi(s + 1)$. The Bethe Ansatz equations (3) for N particles and M down-spins simplify considerably at this point and are in fact the equations for N particles and $M - 1$ down-spins at zero flux. This then gives $D = \chi^{-1}/4\pi^2(s + 1)^2$. Thus we may express the susceptibility in terms of the spin wave velocity as $\chi^{-1} = 2\pi v_s(s + 1)$. In Fig.(1) we show a plot of χ^{-1} as function of the interaction strength. The spin wave velocity has been calculated from the excitation spectrum of the SC model as given in Ref. [6]. As $s \rightarrow 0^-$, the spin wave velocity approaches the velocity of a non-interacting single-component model, i.e. $v_s \rightarrow \pi d/2$. Thus at half-filling, $\chi^{-1}(0^-)$ approaches the non-zero value $\pi^2/2$ which is compatible with the result of Ref. [10].

Finally, we wish to emphasize that we have derived the *exact* Luttinger liquid relation in the SC model starting from the *asymptotic* Bethe Ansatz equations. This further supports the use of the asymptotic Bethe Ansatz as an exact method in the thermodynamic limit.

R.A.R. gratefully acknowledges partial support by the Germanistic Society of America.

REFERENCES

- [1] F. D. M. Haldane, Phys. Rev. Lett. **45**, 1358 (1980); *ibid.* **47**, 1840 (1980); Phys. Letters **81** A, 153 (1981); J. Phys. C **14**, 2585 (1981).
- [2] A. M. Polyakov, Zh. Eksp. Teor. Fiz. Pis. Red. **12**, 538 (1970) [Sov. Phys. JETP Lett. **12**, 381 (1970)]; A. A. Belavin, A. M. Polyakov, A. B. Zamolodchikov, Nuc. Phys. B **241**, 333 (1984); D. Friedan, Z. Qui, S. Shenker, Phys. Rev. Lett. **52**, 1575 (1984); J. L. Cardy, J. Phys. A **20**, L891 (1987), J. L. Cardy, *Conformal Invariance in Phase Transitions and Critical Phenomena*, eds. C. Domb, J. L. Lebowitz, vol. **11**, Academic Press, 1987.
- [3] J. L. Cardy, Nuc. Phys. **B270** [FS16], 186 (1986); I. Affleck, Phys. Rev. Lett. **56**, 746 (1986); H. W. J. Blöte, J. L. Cardy, M. P. Nightingale, *ibid.* **56**, 742 (1986).
- [4] V. E. Korepin, N. M. Bogoliubov, A. G. Izergin, *Quantum Inverse Scattering Method and Correlation Functions*, Cambridge University Press (Cambridge), 1993.
- [5] N. Kawakami, S.-K. Yang, Phys. Rev. Lett. **67**, 2493 (1991); R. A. Römer and B. Sutherland, Phys. Rev. B **48**, 6058 (1993).
- [6] B. Sutherland and R. A. Römer, Phys. Rev. Lett. **71**, 2789 (1993).
- [7] R. A. Römer and B. Sutherland, to be published in Phys. Rev. B **49** (1994).
- [8] C. N. Yang and C. P. Yang, Phys. Rev. **147**, 303 (1966); *ibid.* **150**, 321 (1966); *ibid.* **150**, 321 (1966); *ibid.* **151**, 258 (1966).
- [9] R. A. Römer and B. Sutherland, submitted to Phys. Rev. Lett.
- [10] N. Beyers and C. N. Yang, Phys. Rev. Lett. **7**, 46 (1961); B. Sriram Shastry and B. Sutherland, *ibid.* **65**, 243 (1990); B. Sutherland and B. Sriram Shastry, *ibid.* **65**, 1833 (1990); N. Yu and Michael Fowler, University of Virginia preprint.
- [11] C. N. Yang and C. P. Yang, J. Math. Phys. **10**, 1115 (1969).

FIGURES

FIG. 1. The inverse susceptibility χ^{-1} for the SC model from the Luttinger relation for $-1 < s \leq 0$.